

LASER-ABLATED JET COOLED ROTATIONAL SPECTRA OF LARGE MOLECULES: TESTOSTERONE AND BARBARALONES

IKER LEÓN, *Grupo de Espectroscopia Molecular, Lab. de Espectroscopia y Bioespectroscopia, Unidad Asociada CSIC, Universidad de Valladolid, Valladolid, Spain*; ELENA R. ALONSO^a, *Departamento de Química Física, Universidad del País Vasco (UPV-EHU), Bilbao, Spain*; MAURO MATO, *Department of Analytical and Organic Chemistry, Universidad Rovira i Virgili, Tarragona, Spain*; MIGUEL SANZ NOVO^b, SANTIAGO MATA, *Grupo de Espectroscopia Molecular, Lab. de Espectroscopia y Bioespectroscopia, Unidad Asociada CSIC, Universidad de Valladolid, Valladolid, Spain*; ANTONIO M ECHAVARREN, *Department of Analytical and Organic Chemistry, Universidad Rovira i Virgili, Tarragona, Spain*; JOSÉ L. ALONSO, *Grupo de Espectroscopia Molecular, Lab. de Espectroscopia y Bioespectroscopia, Unidad Asociada CSIC, Universidad de Valladolid, Valladolid, Spain*.

For most people, the name 17 β -Hydroxyandrost-4-en-3-one probably means nothing. When its common name “testosterone” is used instead, one of the most important steroids comes to one’s mind. We have vaporized the solid testosterone by laser ablation, and one conformer has been characterized using chirped-pulse Fourier transform microwave LA-CP-FTMW spectroscopy. The detection of such a large biomolecule highlights the actual sensitivity of rotational spectroscopy combined with laser ablation to tackle large molecules with high melting points. To confirm such an improvement, we will also report the results of 1-(4-tert-Butyl)phenyl)tricyclo[3.3.1.0^{2,8}]nona-3,6-dien-9-one, a large fluxional molecule that is central to the understanding of the phenomena of valence tautomerism.

^aE.R.A. was supported by the Fundación Biofísica Bizkaia (Spain).

^bM.S.N. acknowledges funding from the Spanish “Ministerio de Ciencia, Innovación y Universidades” under predoctoral FPU Grant (FPU17/02987).